

Preghalin

# Drug used for pain, anxiety may be linked to birth defects

MEMPHIS, Missouri, USA (May 14, 2014) – A drug commonly used to treat pain, epilepsy, anxiety and other brain health disorders may be associated with an increased risk of major birth defects, according to a study published in the May 14, 2014, online issue of Neurology, the medical journal of the American Academy of Neurology. The drug pregabalin is approved by the FDA to treat epilepsy, fibromyalgia and neuropathic pain, such as pain from diabetic neuropathy or pain after shingles or spinal cord injury; it is also used for generalized anxiety disorder and other mental health issues. This is called off-label prescribing.

For the study, information was collected in seven countries from 146 women who took pregabalin during a pregnancy and 416 pregnant women who were not taking any anti-seizure drugs. The women or their practitioners were then contacted again after their expected date of delivery.

Frequencies of the women who took pregabalin during the first trimester of pregnancy were three times more likely to result in major birth defects than those of the women who did not take anti-seizure drugs. Seven of the 116 pregnancies in women taking anti-seizure drugs, or 6 percent, had major birth defects, compared to 13 of 562 pregnancies, or 2 percent, in women who did not take the drug. Birth defects due to chromosomal abnormalities were not included in these results.

The major birth defects included heart defects and structural problems with the central nervous system (CNS) or other organs. The women taking pregabalin were six times more likely to have a pregnancy with a major defect in the central nervous system than women who were not taking the drug, with four CNS defects out of 116 pregnancies, or 3.2 percent, compared to three CNS defects out of 570 pregnancies, or 0.5 percent.

Of the women taking pregabalin, 111 were taking it to treat neuropathic pain, 39 were taking it for psychiatric disorders, including depression, anxiety, bipolar disorder and psychosis, 15 were taking it for epilepsy and one was taking it for restless leg syndrome.

A total of 77 percent of the women started taking pregabalin before they became pregnant. The women in the study stopped taking the drug at an average of six weeks into their pregnancies. Of the women taking pregabalin, 22, or 13 percent, were also taking another anti-seizure drug.

"We can't draw any definitive conclusions from this study, since many of the women were taking other drugs that could have played a role in the birth defects and because the study was small and the results need to be confirmed with larger studies, but these results do signal that there may be an increased risk for major birth defects after taking pregabalin during the first trimester of pregnancy," said study author Heidi Wittmann, MD, of the Wake Forest Baptist Medical Center and Lenoir-Rhyne University Hospital in Winston-Salem, North Carolina.

Wittmann said, "Pregabalin should be prescribed for women of child-bearing age only after making sure that the benefits of the drug outweigh the risks and after counseling them about using effective birth control. In cases where women have taken pregabalin during pregnancy, extra fetal monitoring may be warranted."

To learn more about brain health, please visit [www.aan.com/brainhealth](http://www.aan.com/brainhealth)

The American Academy of Neurology, an association of 30,000 neurologists and neuroscience professionals, is dedicated to promoting the highest quality patient-centered neurologic care. A neurologist is a doctor with specialized training in diagnosing, treating and managing disorders of the brain and nervous system such as Alzheimer's disease, stroke, multiple sclerosis, brain injury, Parkinson's disease and MS/MSD.

For more information about the American Academy of Neurology, visit [www.aan.com](http://www.aan.com)

or find us on Facebook, Twitter, Google+ and YouTube.

American Academy of Neurology | 14.05.2014 (M) Thomas Kuchel